

Natural Entanglement in Bose-Einstein Condensates

Christoph Simon

Centre for Quantum Computation, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom
(February 1, 2008)

Every Bose-Einstein condensate is in a highly entangled state, as a consequence of the fact that the particles in a condensate are distributed over space in a coherent way. It is proved that any two regions within a condensate of finite particle number are entangled. This entanglement does not depend on the distance between the two regions. Criteria for the presence of entanglement are derived in the context of interference experiments. For separable states there is a trade-off between fluctuations in particle number and interference visibility.

Entanglement is one of the dramatic non-classical features of quantum physics. There are states of composite quantum systems that cannot be decomposed into probabilistic combinations of product states; there is no way of writing the density matrix of such a state ρ in the form $\rho = \sum_i p_i \sigma_1^i \otimes \sigma_2^i \otimes \dots \otimes \sigma_n^i$, where σ_k^i is a state of the k -th subsystem and the $p_i > 0$ are probabilities. Such states ρ are called inseparable or entangled. The properties of an entangled system are not completely determined by the properties of its parts. Only entangled states can exhibit quantum non-locality [1]. In this case there is no way of reproducing the predictions of quantum physics with classical systems, unless there is instantaneous communication over arbitrary distances.

In the last few years, entanglement has been studied extensively in the context of quantum computation and quantum communication [2]. Methods for creating entangled states have been found in various physical systems, including non-linear optics, ion traps, cavity quantum electrodynamics and nuclear magnetic resonance [3]. A fascinating open problem is whether entanglement plays a significant role in natural physical systems. First steps towards addressing this question were made in the study of interacting spin systems [4]. The effect of decoherence on large-scale spin entanglement has also recently been investigated in this context [5].

Bose-Einstein condensation is another genuine quantum phenomenon which has recently attracted a large amount of experimental and theoretical attention [6]. In a Bose-Einstein condensate (BEC), all particles (which have to be bosons) are in the same quantum state.

Here I would like to draw attention to the fact that by its nature every BEC is a highly entangled state. Even when there is no entanglement of the internal states [7], there is always entanglement in the external degrees of freedom. The discussion will be focussed on the entanglement between different spatial modes, but analogous statements can be made for momentum modes. It will be

shown that any two regions inside a BEC of finite particle number are entangled with each other. The detection of this entanglement via interference will also be discussed.

It is convenient to use the language of non-relativistic quantum field theory. The particles composing the BEC are described by a bosonic quantum field operator $\hat{\psi}(x)$ satisfying the commutation relation $[\hat{\psi}(x), \hat{\psi}^\dagger(y)] = \delta^{(3)}(x - y)\hat{1}$, where x and y are points in three-dimensional space. The time dependence of $\hat{\psi}(x)$ is not essential for the present purpose. Throughout the paper operators will be distinguished by hats.

A BEC of N particles is described by the state $|\Psi\rangle = \frac{1}{\sqrt{N!}}(\hat{f}^\dagger)^N|0\rangle$, where $|0\rangle$ is the vacuum state and the creation operator $\hat{f}^\dagger$ can be written in terms of $\hat{\psi}(x)^\dagger$ as $\hat{f}^\dagger = \int d^3x f^*(x)\hat{\psi}^\dagger(x)$. The mode function $f(x)$ is normalized: $\int d^3x |f(x)|^2 = 1$, where the integrals are over all of space. It is easy to check that $\hat{f}$ fulfills the commutation relation $[\hat{f}, \hat{f}^\dagger] = \hat{1}$.

Let us consider a certain volume inside the BEC and study its entanglement with the rest. Denoting the volume by A , one can define mode operators

$$\begin{aligned}\hat{a} &= \frac{1}{\sqrt{p}} \int_{x \in A} d^3x f(x) \hat{\psi}(x) \\ \hat{b} &= \frac{1}{\sqrt{q}} \int_{x \in \bar{A}} d^3x f(x) \hat{\psi}(x),\end{aligned}\tag{1}$$

where $\bar{A}$ is the complement of A , $p = \int_{x \in A} d^3x |f(x)|^2$ and $q = 1 - p = \int_{x \in \bar{A}} d^3x |f(x)|^2$. Then one has $\hat{f} = \sqrt{p}\hat{a} + \sqrt{q}\hat{b}$, and $\hat{a}$ and $\hat{b}$ fulfill the following commutation relations: $[\hat{a}, \hat{b}] = [\hat{a}, \hat{b}^\dagger] = 0$, $[\hat{a}, \hat{a}^\dagger] = [\hat{b}, \hat{b}^\dagger] = \hat{1}$.

The state of the BEC can now be written as

$$\begin{aligned}|\Psi\rangle &= \frac{1}{\sqrt{N!}}(\sqrt{p}\hat{a}^\dagger + \sqrt{q}\hat{b}^\dagger)^N|0\rangle \\ &= \sum_{k=0}^N \sqrt{\binom{N}{k}} p^{\frac{k}{2}} q^{\frac{N-k}{2}} |k\rangle_a |N-k\rangle_b,\end{aligned}\tag{2}$$

which clearly shows the entanglement between spatial modes $\hat{a}$ and $\hat{b}$. These modes can have different physical meanings. For example, $\hat{a}$ could be the interior of the BEC and $\hat{b}$ the rest of the universe including the BEC's boundary region; $\hat{a}$ and $\hat{b}$ could also each contain one half of the BEC.

The nature of the entanglement in (2) is easy to see. The N particles can be distributed in various ways over

the two modes $\hat{a}$ and $\hat{b}$. The state of the BEC is a coherent superposition of all these possibilities. This clearly generalizes to more than two modes. Note that the particle numbers in $\hat{a}$ and $\hat{b}$ have a binomial distribution.

The present kind of entanglement cannot be detected by local measurements, i.e. measurements acting on modes $\hat{a}$ or $\hat{b}$ separately. If one is restricted to local measurements in the basis of particle number, then (2) is indistinguishable from a separable mixture of the various terms without any fixed phase relation between them. One would therefore require local measurements in bases of states corresponding to superpositions of different particle numbers, which is impossible. However, this does not mean that the entanglement in (2) is unobservable. It can be detected by *joint* measurements on the two modes, for example by interference experiments. This point will be discussed in more detail below.

Let us now study the entanglement properties of BECs in more detail. First suppose that instead of a Fock state with exactly N particles the BEC is in a coherent state of the form [8]

$$|\alpha\rangle = e^{-|\alpha|^2/2} e^{\alpha \hat{f}^\dagger} |0\rangle = e^{-|\alpha|^2/2} e^{\alpha \int d^3x f^*(x) \hat{\psi}^\dagger(x)} |0\rangle. \quad (3)$$

This can formally be rewritten as

$$|\alpha\rangle = e^{-|\alpha|^2/2} \prod_{x \in R^3} e^{\alpha f^*(x) \hat{\psi}^\dagger(x)} |0\rangle, \quad (4)$$

where the product is over all points in space. The state (4) is clearly a product state with respect to the point-like spatial quasi-modes $\hat{\psi}(x)$, since the vacuum does not contain any entanglement [9].

As a consequence all states that can be written as convex combinations of coherent states are also unentangled. In particular this applies to a Poisson distribution of number states $|n\rangle = \frac{(\hat{f}^\dagger)^n}{\sqrt{n!}} |0\rangle$:

$$\rho_P = \sum_{n=0}^{\infty} \frac{e^{-\lambda} \lambda^n}{n!} |n\rangle \langle n|, \quad (5)$$

which can be rewritten as a mixture of coherent states with fixed amplitude $\sqrt{\lambda}$ but random phase

$$\rho_P = \int_0^{2\pi} d\phi |\sqrt{\lambda} e^{i\phi}\rangle \langle \sqrt{\lambda} e^{i\phi}|, \quad (6)$$

as can easily be shown by expanding the coherent states in the number state basis: $|\alpha\rangle = e^{-|\alpha|^2/2} \sum \frac{\alpha^n}{\sqrt{n!}} |n\rangle$ and performing the integration over ϕ .

Note that a state with a thermal distribution of particle numbers is also not entangled, because it can be written as a convex combination of coherent states with a probability distribution that is a Gaussian in the coherent state amplitude:

$$\rho_T = (1-t) \sum_{n=0}^{\infty} t^n |n\rangle \langle n| = \int d^2\alpha e^{-\beta|\alpha|^2} |\alpha\rangle \langle \alpha|, \quad (7)$$

where the integration is over the complex plane and $\beta = (1/t) - 1$. Again the identity can be shown by expressing the coherent states in the number state basis and performing the integration.

The above results suggest that the fixedness of the particle number plays an important role in ensuring that the state $\frac{(\hat{f}^\dagger)^N}{\sqrt{N!}} |0\rangle$ is entangled. This leads naturally to the question of subsystems. If one considers a volume containing just part of a BEC, then the particle number in this volume is no longer fixed. Does the reduced state corresponding to such a volume still exhibit spatial entanglement of the above type? It will now be shown that the answer is yes, and that the entanglement disappears only in the limit of a finite subsystem of an infinitely large BEC.

Let us partition space into three regions, A , B and C . For concreteness, imagine A and B to be well within the region occupied by the BEC. A and B may be adjacent or separated. We will trace over the region C and study the entanglement between the regions A and B . Defining modes $\hat{a}$, $\hat{b}$ and $\hat{c}$ corresponding to the three regions A , B and C as above, we have $\hat{f}^\dagger = \sqrt{p} \hat{a}^\dagger + \sqrt{q} \hat{b}^\dagger + \sqrt{1-p-q} \hat{c}^\dagger$, where p , q and $1-p-q$ are the probabilities for an individual particle to be found in regions A , B and C respectively. We now trace the state $|\Psi\rangle = \frac{(\hat{f}^\dagger)^N}{\sqrt{N!}} |0\rangle$ over the $\hat{c}$ -mode in order to find the reduced density matrix for modes $\hat{a}$ and $\hat{b}$. Applying Eq. (2) to the split $AB-C$ it is clear that the reduced density matrix will correspond to the states

$$\frac{(\sqrt{\frac{p}{p+q}} \hat{a}^\dagger + \sqrt{\frac{q}{p+q}} \hat{b}^\dagger)^n}{\sqrt{n!}} |0\rangle \quad (8)$$

occurring with binomial probabilities $\binom{N}{n} (p+q)^n (1-p-q)^{N-n}$.

First consider the limit where A and B are finite-size subsystems of an infinitely large BEC. The total particle number N goes to infinity in such a way that the mean particle number in systems A and B together, $\lambda = N(p+q)$, remains finite. It is easy to show that in this limit the binomial distribution tends towards a Poisson distribution. As a consequence the reduced state is exactly of the form (5) with the individual number states of the form (8). Thus in this limit there is no entanglement in the system consisting of A and B only.

As long as C is of finite size, there is always entanglement between A and B . To show this, let us write the reduced state of modes $\hat{a}$ and $\hat{b}$ explicitly:

$$\text{Tr}_C |\Psi\rangle \langle \Psi| = \sum_{n=0}^N \sum_{k,l=0}^n \binom{N}{n} (1-p-q)^{N-n} \sqrt{\binom{n}{k} \binom{n}{l}} (\sqrt{p})^{k+l} (\sqrt{q})^{2n-k-l} |k, n-k\rangle \langle l, n-l|, \quad (9)$$

where $|k, n-k\rangle = \frac{(\hat{a}^\dagger)^k (\hat{b}^\dagger)^{n-k}}{\sqrt{k!(n-k)!}}|0\rangle$. The inseparability of (9) can be shown using the Peres-Horodecki partial transposition criterion [10].

This can be done by first projecting the state onto suitable two-dimensional subspaces in both modes. The simplest choice are the subspaces spanned by the states $|0\rangle$ and $|1\rangle$. The resulting state is

$$(1-p-q)^N|00\rangle\langle 00| + N(1-p-q)^{N-1}(p|10\rangle\langle 10| + q|01\rangle\langle 01| + \sqrt{pq}|10\rangle\langle 01| + \sqrt{pq}|01\rangle\langle 10|) + \frac{N(N-1)}{2}(1-p-q)^{N-2}2pq|11\rangle\langle 11|, \quad (10)$$

whose partial transpose has a negative eigenvalue for all values of p and q , which is seen most easily by calculating its determinant. Thus any two regions within a finite-size BEC are entangled with each other.

This is not the only possible choice of two-dimensional subspaces which exhibits entanglement. If one projects in both modes onto two arbitrary Fock states $|n\rangle$ and $|m\rangle$ (the same two numbers for modes $\hat{a}$ and $\hat{b}$), the resulting density matrix also has a non-positive partial transpose for all values of p and q , which can be shown in the same way as before. The amount of entanglement between the two regions A and B depends only on the total particle number N and on the average particle numbers in the two regions, determined by p and q . It does not depend on the distance between the two regions. This emphasizes the spatially coherent character of the BEC.

We have noted above that the present type of entanglement cannot be detected by separate measurements on the modes $\hat{a}$ and $\hat{b}$. However it can be detected by joint measurements of the two modes. A (conceptually) simple possibility is to perform an interference experiment. For simplicity consider the state

$$|\psi\rangle = \frac{(\hat{a}^\dagger + \hat{b}^\dagger)^N}{\sqrt{2^N N!}}|0\rangle, \quad (11)$$

which has $\langle\psi|\hat{a}^\dagger\hat{a}|\psi\rangle = \langle\psi|\hat{b}^\dagger\hat{b}|\psi\rangle = N/2$ and a fixed total particle number N . Now consider detection in the new basis of modes given by $\hat{a}' = (\hat{a} + \hat{b})/\sqrt{2}$, $\hat{b}' = (\hat{a} - \hat{b})/\sqrt{2}$. This corresponds to a measurement after superimposing modes $\hat{a}$ and $\hat{b}$ on a 50-50 beam splitter. The state (11) satisfies $\langle\psi|\hat{a}'^\dagger\hat{a}'|\psi\rangle = N$ and $\langle\psi|\hat{b}'^\dagger\hat{b}'|\psi\rangle = 0$. As a consequence of interference, behind the beam splitter all particles are concentrated in one of the modes.

There is no separable state of modes $\hat{a}$ and $\hat{b}$ which has the same properties, namely (i) a fixed total particle number (different from zero), i.e. $\langle\hat{N}\rangle^2 = \langle\hat{N}^2\rangle - \langle\hat{N}\rangle^2 = 0$, where $\hat{N} = \hat{a}^\dagger\hat{a} + \hat{b}^\dagger\hat{b} = \hat{a}'^\dagger\hat{a}' + \hat{b}'^\dagger\hat{b}'$, and (ii) complete destructive interference, i.e. $\langle\hat{b}'^\dagger\hat{b}'\rangle = 0$, at the same time.

This can be seen in the following way. Consider a general separable state of modes $\hat{a}$ and $\hat{b}$:

$$\rho = \sum p_i |\phi_i\rangle\langle\phi_i| \otimes |\chi_i\rangle\langle\chi_i|, \quad (12)$$

where $|\phi_i\rangle$ are states of mode $\hat{a}$ and $|\chi_i\rangle$ are states of mode $\hat{b}$.

First assume that the particle number is fixed to be N . This implies $\hat{N}|\phi_i\rangle|\chi_i\rangle = N|\phi_i\rangle|\chi_i\rangle$ for all i . This is possible only if $|\phi_i\rangle$ is an eigenstate of $\hat{a}^\dagger\hat{a}$ and $|\chi_i\rangle$ is an eigenstate of $\hat{b}^\dagger\hat{b}$, i.e. $|\phi_i\rangle|\chi_i\rangle = |n_i\rangle|N-n_i\rangle$. So ρ must be a convex combination of products of Fock states. But Fock states have no definite phase, therefore destructive interference cannot occur. It is easy to see that in this case one has $\langle\hat{b}'^\dagger\hat{b}'\rangle = \text{Tr}\rho\hat{b}'^\dagger\hat{b}' = N/2$. The particles have a 50-50 distribution behind the beam splitter, in contrast to the result for (11).

On the other hand, one can also impose $\langle\hat{b}'^\dagger\hat{b}'\rangle = 0$. This implies

$$\frac{1}{2} \sum p_i \left(\langle\hat{a}^\dagger\hat{a}\rangle_i + \langle\hat{b}^\dagger\hat{b}\rangle_i - \langle\hat{a}^\dagger\rangle_i\langle\hat{b}\rangle_i - \langle\hat{a}\rangle_i\langle\hat{b}^\dagger\rangle_i \right) = 0, \quad (13)$$

where we have introduced the shorthand notation $\langle\hat{a}^\dagger\hat{a}\rangle_i = \langle\phi_i|\hat{a}^\dagger\hat{a}|\phi_i\rangle$, $\langle\hat{b}^\dagger\hat{b}\rangle_i = \langle\chi_i|\hat{b}^\dagger\hat{b}|\chi_i\rangle$, $\langle\hat{a}\rangle_i = \langle\phi_i|\hat{a}|\phi_i\rangle$, $\langle\hat{b}\rangle_i = \langle\chi_i|\hat{b}|\chi_i\rangle$ etc.

Consider the expression in parentheses under the sum in (13), dropping the index i for a moment. If we define $\langle\hat{a}\rangle = Ae^{i\alpha}$ and $\langle\hat{b}\rangle = Be^{i\beta}$ with A, B positive and α, β real, this translates into

$$A^2 + C_a + B^2 + C_b - 2AB\cos(\alpha - \beta). \quad (14)$$

Here we have introduced the correlation functions $C_a = \langle\hat{a}^\dagger\hat{a}\rangle - \langle\hat{a}\rangle\langle\hat{a}^\dagger\rangle$ and $C_b = \langle\hat{b}^\dagger\hat{b}\rangle - \langle\hat{b}\rangle\langle\hat{b}^\dagger\rangle$, which are positive definite, as can be seen from the Cauchy-Schwarz inequality applied to vectors $|\phi\rangle$ and $|v\rangle = \hat{a}|\phi\rangle$ [11].

Therefore (14) is never negative, and equal to zero only if $C_a = C_b = 0$, $A = B$ and $\cos(\alpha - \beta) = 1$. The Cauchy-Schwarz inequality is saturated only if the two vectors are collinear, therefore the first condition implies $\hat{a}|\phi\rangle = Ae^{i\alpha}|\phi\rangle$ and $\hat{b}|\phi\rangle = Be^{i\beta}|\phi\rangle$. Returning to Eq. (13), this implies that all the states $|\phi_i\rangle$ and $|\chi_i\rangle$ have to be coherent states, with the additional constraints $A_i = B_i$ and $\cos(\alpha_i - \beta_i) = 1$ for all i . This implies that the particle number cannot be fixed. For a mixture of products of coherent states of the form $\sum_i p_i |A_i\rangle\langle A_i| \otimes |A_i\rangle\langle A_i|$ one has $\langle\hat{N}\rangle = \langle\hat{a}^\dagger\hat{a} + \hat{b}^\dagger\hat{b}\rangle = 2\sum_i p_i |A_i|^2$ and $\langle\hat{N}^2\rangle = \langle\hat{a}^\dagger\hat{a}\hat{a}^\dagger\hat{a} + \hat{b}^\dagger\hat{b}\hat{b}^\dagger\hat{b} + 2\hat{a}^\dagger\hat{a}\hat{b}^\dagger\hat{b}\rangle = 4\sum_i p_i |A_i|^4 + 2\sum_i p_i |A_i|^2$, using $\hat{a}\hat{a}^\dagger = \hat{a}^\dagger\hat{a} + \hat{1}$. Since $\sum_i p_i |A_i|^4$ is always greater than or equal to $(\sum_i p_i |A_i|^2)^2$, this implies that $(\Delta\hat{N})^2 = \langle\hat{N}^2\rangle - \langle\hat{N}\rangle^2 \geq \langle\hat{N}\rangle$. If one demands perfect destructive interference, then (for a separable state) the particle number cannot be definite.

Thus for separable states there is clearly a trade-off between the fluctuations in particle number, expressed by $(\Delta\hat{N})^2$, and the amount of interference, expressed by $\langle\hat{b}'^\dagger\hat{b}'\rangle$. If one of the two quantities is zero, the other one

is bounded away from zero. If these lower bounds are violated, the state under consideration must be entangled. It may be possible to extend the above results, for example to derive a region in the $((\Delta\hat{N})^2, \langle\hat{b}^\dagger\hat{b}'\rangle)$ plane (for fixed mean particle number) where the states are definitely entangled. This would be desirable because in a real experiment neither of the two quantities would be exactly equal to zero. Such a more general criterion might then allow a direct experimental proof of the presence of entanglement in Bose-Einstein condensates via interference. Note that interference between different distinct regions of a Bose-Einstein condensate was observed in [12], where the BEC was distributed over an array of micro-traps.

The effect of non-zero temperature on the entanglement analyzed here is another topic for future research. More generally, one may wonder whether new insights in condensate phenomena such as superfluidity can be gained from the point of view of entanglement. Finally it should be interesting to compare the present results to the case of condensed fermion pairs.

I would like to thank S. Bose and A. Costa for useful discussions. This work was supported by the QuComm project of the European Union (IST-1999-10033).

ber superselection rule, which may be a fundamental law of nature or due to very fast environmentally induced decoherence. For a discussion of superselection rules cf. D. Giulini, quant-ph/0010090. Here these states are used only as a convenient mathematical tool for analyzing the properties of the mixed states (5) and (7).

- [9] Our present discussion is within the framework of non-relativistic quantum field theory. It is worth noting that in relativistic quantum field theory even the vacuum is an entangled state. See e.g. S.J. Summers and R. Werner, Phys. Lett. A **110**, 257 (1985); L.J. Landau, Phys. Lett. A **123**, 115 (1987); H. Halvorson and R. Clifton, J. Math. Phys. **41**, 1711 (2000); B. Reznik, quant-ph/0008006 (2000).
- [10] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996); M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 1 (1996).
- [11] $\langle v|v\rangle = \langle\phi|\hat{a}^\dagger\hat{a}|\phi\rangle$ and $\langle\phi|v\rangle = \langle\phi|\hat{a}|\phi\rangle$ fulfill the Cauchy-Schwarz inequality $\langle v|v\rangle\langle\phi|\phi\rangle \geq \langle v|\phi\rangle\langle\phi|v\rangle$.
- [12] B.P. Anderson and M. Kasevich, Science **282**, 1686 (1998).

-
- [1] J.S. Bell, Physics **1**, 195 (1964); J.S. Bell, *Speakable and Unsayable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987).
 - [2] M.A. Nielsen and I.L. Chuang, *Quantum computation and quantum information* (Cambridge University Press, Cambridge, 2000).
 - [3] D. Bouwmeester, A. Ekert, and A. Zeilinger, *The Physics of Quantum Information* (Springer, Berlin, 2000).
 - [4] M.C. Arnesen, S. Bose, and V. Vedral, Phys. Rev. Lett. **87**, 017901 (2001); K.M. O'Connor and W.K. Wootters, Phys. Rev. A **63**, 052302 (2001); W.K. Wootters, quant-ph/0001114; X. Wang, Phys. Rev. A **64**, 012313 (2001); X. Wang, Phys. Lett. A **281**, 101 (2001); D. Gunlycke, V.M. Kendon, V. Vedral, and S. Bose, Phys. Rev. A **64**, 042302 (2001).
 - [5] C. Simon and J. Kempe, quant-ph/0109102 (2001).
 - [6] M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman, and E.A. Cornell, Science **269**, 198 (1995); K.B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995); A.S. Parkins and D.F. Walls, Phys. Rep. **303**, 1 (1998).
 - [7] A. Sørensen, L.-M. Duan, J.I. Cirac, and P. Zoller, Nature **409**, 63 (2001) and K. Helmerson and L. You, Phys. Rev. Lett. **87**, 170402 (2001) propose ways of creating entanglement between the internal degrees of freedom of particles in a BEC.
 - [8] Such superposition states of different particle numbers are impossible to prepare in practice. If the particles are atoms or molecules, such states violate the baryon num-